Journal of Pharmaceutical and Biomedical Analysis

Volumes 4 and 5, 1986–1987

LIST OF CONTENTS

Volume 4

Number 1

C. A. Johnson — 1 — Editorial

Analytical Survey

R. D. McDowall, J. C. Pearce and G. S. Murkitt — 3 — Liquid–solid sample preparation in drug analysis

Full Papers

L. L. Lewin, P. Sanz Pedrero and M. R. Vilarino — 23 — Direct determination of zinc in insulin preparations by differential pulse polarography

J. H. Gill and G. M. Holder — 31 — Application of synchronous luminescence to the separate determination of cochromatographing metabolites of the carcinogen, 7-methylbenz[*c*]acridine

P.-A. Johansson, G. Östling and J. Drakenberg — 37 — Rapid spectrophotometric pH measurements based on the dual-wavelength technique

V. V. Coşofreţ and R. P. Buck — 45 — A poly(vinylchloride) membrane electrode for determination of phenytoin in pharmaceutical formulations

B. Rollman, P. Van der Auwera and P. M. Tulkens — 53 — Dibekacin assay in serum by automated fluorescence polarization immunoassay (Abbott Tdx): Comparison with high-performance liquid chromatography, substrate-labelled fluorescent immunoassay and radioimmunoassay

K.-O. Eriksson and S. Hjertén — 63 — Estimation of peptide/protein molecular weights by high-performance molecular-sieve chromatography on agarose columns in 6 M guanidine hydrochloride

S. H. Hansen and S. B. Pedersen — 69 — Assay of tolfenamic acid and its metabolites by liquid chromatography on dynamically modified silica: application in pharmacokinetics

C. M. Selavka, I. S. Krull and K. Bratin — 83 — Analysis for penicillins and cefoperazone by HPLC-photolysis-electrochemical detection (HPLC-hv-EC)

M. Hassan and H. Ehrsson — 95 — Degradation of busulfan in aqueous solution

Short Communications

J. S. O'Neal, K. B. Sloan and S. G. Schulman — 103 — Substrate-labelled fluorescence immunoassay of phenytoin

H. Ueda and J. H. Perrin — 107 — A microcalorimetric investigation of the binding of flurbiprofen to cyclodextrins

J. R. Andersen and P. Helboe — 111 — Aluminium in antihaemophilia preparations as determined by electrothermal atomic absorption spectrophotometry

J. C. Pearce, G. S. Murkitt, D. C. Taylor and P. R. Cresswell — 115 — An assay for prizidilol in plasma and urine by high-performance liquid chromatography

G. Szepesi, M. Gazdag, Zs. Pap-Sziklay and Z. Végh — 123 — Problems of semiquantitative TLC methods prescribed in the United States and British Pharmacopoeias for the purity testing of sulphinpyrazone

R. C. Young and R. W. Nysewander — 131 — Changes in plasma and erythrocyte chlorpromazine concentrations during sample storage

Number 2

A. F. Fell and J. N. Miller — 135 — Editorial

Analytical Survey

J. C. Lindon — 137 — NMR spectroscopy: analytical applications from chemistry to the clinic

Full Papers

S. T. Eberhart, A. Hatzis and R. Rothchild — 147 — Quantitative NMR assay for aspirin, phenacetin, and caffeine mixtures with 1,3,5-trioxane as internal standard

L. Campanella, M. Tomassetti and M. Cordatore — 155 — Application of a new cholate liquid membrane electrode to the determination of the cholic acids pool in human bile

P. Demedts, J. Van der Verren and A. Heyndrickx — 163 — The conversion of a standard gas chromatograph to a fused silica capillary system: applications in human toxicology and tropical medicine

I. M. Johansson, E.-L. Anlér, U. Bondesson and B. Schubert — 171 — Isolation of meclofenamic acid and two metabolites from equine urine—a comparison between horse and man

L.-E. Edholm — 181 — Coupled columns in bioanalytical work

D. Stevenson, R. Briggs and G. P. Mould — 191 — A bioavailability study of two preparations of tamoxifen after single doses

H. Derendorf, P. Rohdewald, G. Hochhaus and H. Möllmann — 197 — HPLC determination of glucocorticoid alcohols, their phosphates and hydrocortisone in aqueous solutions and biological fluids

H. J. Ballard, D. Cotterrell and F. Karim — 207 — Analysis of submicromolar concentrations of adenosine in plasma using reversed phase high-performance liquid chromatography

C. Pettersson, T. Arvidsson, A.-L. Karlsson and I. Marle — 221 — Chromatographic resolution of enantiomers using albumin as complexing agent in the mobile phase

C. L. Baustian and T. J. Mikkelson — 237 — Analysis of clomiphene isomers in human plasma and detection of metabolites using reversed-phase chromatography and fluorescence detection

C. L. Willis, C. K. Lim and T. J. Peters — 247 — Separation of ribonucleotides, deoxynucleotides, cyclic nucleotides and deoxycyclic nucleotides by reversed-phase high-performance liquid chromatography

N. Daoud, T. Arvidsson and K.-G. Wahlund — 253 — The liquid chromatographic determination of theophylline in untreated plasma by simple direct injection

N.-O. Ahnfelt, R. Modin, M.-L. Odell and U. Ohlson — 261 — Determination of salicylazosulphapyridine, sulphapyridine and its metabolites in serum and urine

Short Communication

S. E. Patterson — 271 — Determination of temazepam in plasma and urine by high-performance liquid chromatography using disposable solid-phase extraction columns

Number 3

Review

J. H. Beijnen, H. Lingeman, H. A. Van Munster and W. J. M. Underberg — 275 — Mitomycin antitumour agents: a review of their physico-chemical and analytical properties and stability

Full Papers

M. R. Detaevernier, Y. Michotte, L. Buydens, M. P. Derde, M. Desmet, L. Kaufman, G. Musch, J. Smeyers-Verbeke, A. Thielemans, L. Dryon and D. L. Massart — 297 — Feasibility study concerning the use of expert systems for the development of procedures in pharmaceutical analysis

R. Jones and B. S. Leadley — 309 — High pass digital filtering applied to quantitative UV spectrometry

M. M. Andino and J. D. Winefordner — 317 — Room temperature phosphorescence of diazepam and its application to the determination of diazepam in serum and in a tablet formulation

D. S. Skrabalak and J. D. Henion — 327 — Identification of betamethasone and a major metabolite in equine urine

V. Marko, J. Wijsbeek and R. A. de Zeeuw — 333 — Determination of carbisocaine, heptacaine and pentacaine in plasma by capillary gas chromatography with nitrogen-selective detection

J. F. Stobaugh, A. J. Repta and L. A. Sternson — 341 — Aspects of the stability of isoindoles derived from the reaction of *o*-phthalaldehyde–ethanethiol with primary amino compounds

D. M. Maniez-Devos, R. Baurain, M. Lesne and A. Trouet — 353 — Degradation of doxorubicin and daunorubicin in human and rabbit biological fluids

P. Cockaerts, E. Roets and J. Hoogmartens — 367 — Analysis of a complex analgesic formulation by high-performance liquid chromatography with column-switching

B. V. Fisher and D. Smith — 377 — HPLC as a replacement for the animal response assays for insulin

I. Grundevik, G. Jerndal, K. Balmér and B.-A. Persson — 389 — Fully automated gradient elution liquid chromatographic assay of omeprazole and two metabolites

M. Abrahamsson — 399 — Determination of a prodrug of tranexamic acid in whole blood by reversed-phase liquid chromatography after pre-column derivatization with fluorescamine

Short Communications

R. C. Young and R. W. Nysewander — 407 — Plasma concentrations of chlorpromazine and blood collection method

B. de Spiegeleer, G. Slegers and W. van den Bossche — 411 — Rapid determination of malonic acid by ion-pair reversed-phase high-performance liquid chromatography

Number 4

Review

J. Bélanger and J. R. J. Paré — 415 — Fast atom bombardment mass spectrometry in the pharmaceutical analysis of drugs

Full Papers

A. Hatzis and R. Rothchild — 443 — Optical purity determination and ^{1}H-NMR spectral simplification with lanthanide shift reagents—VIII. An indacrinone precursor, 6,7-dichloro-5-methoxy-2-methyl-2-phenyl-1-indanone

A. Hatzis and R. Rothchild — 451 — ^{1}H NMR spectral simplification with achiral and chiral lanthanide shift reagents. Tranylcypromine, *trans*-2-phenylcyclopropanamine

N. A. Sparrow, L. Glasser, D. Glasser and A. E. Russell — 461 — Binding isotherms by continuous-flow dynamic dialysis

S. Zommer-Urbańska and H. Bojarowicz — 475 — Spectrophotometric investigations on protolytic equilibria of mefenamic acid and determination by means of Fe(III) in methanol–aqueous media

J. M. Gallo, C. T. Hung and D. G. Perrier — 483 — Reversed-phase ion-pair high-performance liquid chromatography of adriamycin and adriamycinol in rat serum and tissues

I. A. Eiseman, D. L. Reynolds, E. L. Johnson, A. C. Daftsios and L. A. Pachla — 491 — A validated liquid chromatographic method for the antiallergy agent CI-922 in dog and rat plasma

M. Vajda, Zs. Tegyey, M. Kozma and L. Vereczkey — 497 — Studies on the pharmacokinetics and metabolism of N_4-carbamoyl-1,3,4,5-dihydro-diazepam (Uxepam®) in rats, dogs and man

Short Communications

M. Balón-Almeida, M. A. Muñoz-Perez, J. Hidalgo-Toledo and M. Sánchez-González — 505 — Fluorescence characteristics of Rauwolfia alkaloids in highly concentrated sulphuric acid solutions. Fluorimetric determination of reserpiline

S. K. Talwar, S. C. Sharma and S. Das — 511 — Simultaneous spectrophotometric determination of diloxanide furoate and metronidazole in dosage forms

A. Gergely and Gy. Szász — 517 — Use of circular dichroism spectroscopy for the determination of oily injections containing a Δ^4-3-ketosteroid

J. H. Wolf, C. de Ruiter, U. A. Th. Brinkman and R.W. Frei — 523 — Liquid chromatographic determination of a quaternary ammonium-steroidal type drug in plasma with post-column extraction

H. Omori, Y. Takahashi, T. Sakakibara, S. Ota and T. Nakashizuka — 529 — The determination of bradykinin in human saliva by high-performance liquid chromatography with UV detection

T.-M. Chen, A. D. Sill and C. L. Housmyer — 533 — Solution stability study of terfenadine by high-performance liquid chromatography

S. Linko-Löppönen and E. Nissinen — 541 — Liquid-chromatographic analysis of a novel antitussive agent, 2′,4′-dimethyl-6′-methoxy-3-(2-methyl-piperidyl)propionanilide in human urine

Number 5

Reviews

J. Ruz, A. Fernandez, M. D. L. de Castro and M. Valcarcel — 545 — Determination of ethanol in human fluids—I. Determination of ethanol in blood

J. Ruz, A. Fernandez, M. D. L. de Castro and M. Valcarcel — 559 — Determination of ethanol in human fluids—II. Determination of ethanol in urine, breath and saliva

Analytical Survey

C. A. Johnson — 565 — Purity requirements from a pharmacopoeial point of view

Full Papers

S. A. Botha, J. K. Guillory and A. P. Lötter — 573 — Physical characterization of solid forms of urapidil

S. P. Musial, M. P. Duran and R. V. Smith — 589 — Analysis of solvent-mediated conformational changes of insulin by radioimmunoassay (RIA) techniques

D. J. Mazzo, E. K. F. Fong and S. E. Biffar — 601 — A comparison of test methods for determining *in vitro* drug release from transdermal delivery dosage forms

K. Bergström — 609 — Carbohydrates—purity assessment

S.-O. Jansson — 615 — Characterization of drug purity by liquid chromatography

P. O. Edlund — 625 — Determination of dihydroxyphenylalanine and dihydroxyphenylacetic acid in biological samples by coupled-column liquid chromatography with dual coulometric–amperometric detection

P. O. Edlund — 641 — Automated analysis of homovanillic acid and 5-hydroxyindole-3-acetic acid in biological fluids by coupled-column liquid chromatography and dual coulometric–amperometric detection

J. W. Cox and R. H. Pullen — 653 — Determination of prostaglandin E_1 in plasma with picogram per millilitre sensitivity by double antibody extraction and column switching high-performance liquid chromatography

Short Communications

I. D. Wilson and I. M. Ismail — 663 — A rapid method for the isolation and identification of drug metabolites from human urine using solid phase extraction and proton NMR spectroscopy

B. B. Ba, H. Bun, P. Coassolo and J. P. Cano — 667 — Rapid determination of the metabolic pool of ethyl loflazepate in biological fluids by capillary gas chromatography

V. Rossetti, A. Lombard and M. Buffa — 673 — The HPTLC resolution of the enantiomers of some 2-arylpropionic acid anti-inflammatory drugs

A. Pitotti, V. Maurich, M. Moneghini and S. Vianello — 677 — HPLC method for evaluation of urinary angiotensin-converting enzyme: some examples of normal subjects and patients with renal transplantation

Number 6

Special Issue: Drug Analysis 1986

A. F. Fell and J. N. Miller — 685 — Editorial

Plenary Papers

C. A. Johnson — 689 — Drug analysis—Why?

P. Crevat-Pisano, C. Hariton, P. H. Rolland and J. P. Cano — 697 — Fundamental aspects of radioreceptor assays

J. W. Munson — 717 — Analytical techniques in dissolution testing and bioavailability studies

J. van Rompay — 725 — Purity determination and evaluation of new drug substances

Keynote Papers

A. P. de Leenheer, M. F. Lefevere and L. M. R. Thienpont — 735 — Bioanalytical mass spectrometry as applied to organic acid profiling and steroid reference methodology

C. J. de Ranter — 747 — Applications of X-ray diffractometric techniques in the analysis of drugs

D. Giron — 755 — Applications of thermal analysis in the pharmaceutical industry

G. G. Guilbault — 771 — Use of luminescence spectroscopy for assay of pharmaceutical compounds

J. R. Kirchhoff, E. Deutsch and W. R. Heineman — 777 — Characterization of technetium radiopharmaceuticals by thin-layer spectroelectrochemistry

G. J. Patriarche — 789 — New trends on modified electrodes: applications to drug analysis

G. Szepesi, Zs. Pap-Sziklay, Zs. Bertha-Somodi and Z. Vegh — 799 — Use of overpressurized thin-layer chromatography (OPTLC) for the separation of amino-acids and polypeptides

P. V. Huong — 811 — Drug analysis by Raman and micro-Raman spectroscopy

J. Vessman, K. E. Karlsson and O. Gyllenhaal — 825 — Direct derivatization of drugs in untreated biological samples for gas chromatographic analysis

Volume 5 Number 1

Full Papers

M. A. Iorio, A. M. Farina and A. Doldo — 1 — Quantitative analysis of β-adrenergic blocking agents by NMR spectroscopy

K. Yamashita, K. Watanabe, H. Takayama, M. Ishibashi and H. Miyazaki — 11 — Measurement of plasma etoposide by radioimmunoassay

V. Cavrini, A. M. di Pietra, M. A. Raggi and R. Sarti — 21 — Analysis of nandrolone esters in oily injections by reversed-phase high-performance liquid chromatography

C. M. Riley and A. K. Runyan — 33 — High-performance liquid chromatography of anthracycline antibiotics with electrochemical detection. Application to the clinical pharmacokinetics of 4′-deoxydoxorubicin

Short Communications

G. I Adebayo and A. F. B. Mabadeje — 45 — Better response to thiazides in blacks—lack of pharmacokinetic evidence with chlorothiazide

K. K. Verma, P. Tyagi and A. K. Gulati — 51 — Determination of mixtures of sulphonamides by titration with chloramine-T

M. S. Ionescu, A. A. Abrutis, M. Lăzărescu, E. Pascu, G. E. Baiulescu and V. V. Coşofreţ — 59 — Amantadine ion-selective membrane electrodes and their use in pharmaceutical analysis

E. M. May, D. C. Hunt and J. Sloggem — 65 — A normal-phase high-performance liquid chromatographic assay for aziridine residue in trientine dihydrochloride

71 — Announcement

Number 2

Review

S. K. Branch, A. F. Casy and E. M. A. Ominde — 73 — Application of ^{1}H nuclear magnetic resonance spectroscopy to the analysis of β-lactam antibiotics and their common degradation products

Full Papers

C. J. M. Sindic — 105 — The use of particle counting immunoassay for the diagnosis of neurological disease

P. L. Masson — 113 — Particle counting immunoassay—an overview

A. Hatzis and R. Rothchild — 119 — Optical purity determination, conformer populations and ^{1}H NMR spectral simplification with lanthanide shift reagents—Part IX. A method for improved analytical precision for "DOEt", 2,5-dimethoxy-4-ethylamphetamine

J. R. J. Paré, P. Lafontaine, J. Bélanger, Wing-Wah Sy, N. Jordan and J. C. K. Loo — 131 — Characterization of digoxin and related cardiac glycosides by fast atom bombardment mass spectrometry

K. Karljiković-Rajić, B. Stanković and Z. Binenfeld — 141 — Spectrophotometric determination of obidoxime chloride as its Pd(II) complex in water and injections

R. M. Facino, M. Carini, D. Castoldi and O. Tofanetti — 151 — Analysis of tiadenol in human plasma by capillary gas chromatography with electron capture detection

Short Communications

I. D. Wilson, J. Fromson, I. M. Ismail and J. K. Nicholson — 157 — Proton magnetic resonance spectroscopy of human urine: excretion of 1-(3′-carboxypropyl)-3,7-dimethylxanthine by man after dosing with oxpentifylline

R. A. Baumann, C. Gooijer, N. H. Velthorst, R. W. Frei, I. Klein and W. J. F. van der Vijgh — 165 — Quantitative determination of cisplatin in body fluids by liquid chromatography with quenched phosphorescence detection

C. Bertucci, C. Rosini, D. Pini and P. Salvadori — 171 — Chiral stationary phases and circular dichroism detection in high-performance liquid chromatography: determination of stereochemical purity of drugs

A. Bernareggi, P. Poletti, G. Zanolo and L. F. Zerilli — 177 — Simultaneous determination of the two main metabolites of deflazacort in human plasma by high-performance liquid chromatography

K. Kawahara, T. Awaji, K. Uda and Y. Sakai — 183 — Determination of four metabolites of dothiepin in urine by high-performance liquid chromatography

Number 3

Review
E. Tyihák — 191 — Overpressured layer chromatography and its applicability in pharmaceutical and biomedical analysis

Full Papers
R. Gimet and A. T. Luong — 205 — Quantitative determination of polymorphic forms in a formulation matrix using the near infra-red reflectance analysis technique

A. G. Davidson and T. O. Dawodu — 213 — Difference spectrophotometric assay of 5-hydroxymethylfurfuraldehyde in hydrolysed pharmaceutical syrups—I. Sodium borohydride reagent

F. Mebsout, J.-M. Kauffmann and G. J. Patriarche — 223 — Redox behaviour of anti-tumor platinum (II) compounds (carboplatin) at solid electrodes

S. Thomson Eberhart, A. Hatzis, J. Jimenez, R. Rothchild and P. Simons — 233 — ^{1}H NMR spectral simplification with achiral and chiral lanthanide shift reagents—IV. Thiopental and barbiturate analogues

A. F. Casy — 247 — Electron-impact mass spectrometry of diuretic agents

E. Besenfelder — 259 — The determination of torasemide and metabolites in plasma by high-performance liquid chromatography

J. A. Adamovics — 267 — Determination of antibiotics and antimicrobial agents in human serum by direct injection onto silica liquid chromatographic columns

S. H. Curry, S. M. Aburawi and R. Whelpton — 275 — Analysis of solutions containing glutathione and inorganic nitrite: application to nitroglycerin metabolism studies

Short Communications
C. Franconi, M. Bonori, E. F. Orsega, M. Scarpa and A. Rigo — 283 — Enzyme immunoassay by polarography—A preliminary study

N. Sadlej-Sosnowska — 289 — Determination of glycyrrhizinic and glycyrrhetinic acids in pharmaceuticals by high-performance liquid chromatography

H. W. Jun, N. H. Foda and J. W. McCall — 295 — Reversed-phase liquid chromatographic determination of meclozine in biological samples

G. Mullersman, S. Toufflin and H. Derendorf — 303 — HPLC analysis of buprenorphine in plasma and urine using coulometric detection

Number 4

Review
E. Postaire, F. Pradier, M. Postaire, D. Pradeau, L. Matchoutsky, P. Prognon and M. Hamon — 309 — Various techniques for the routine evaluation of the degradation of glucose in parenteral solutions—a critical study

Analytical Survey
A. H. Thomas — 319 — Replacement of microbiological assay by high-performance liquid chromatographic assay for antibiotics

Full Papers
Shou-zhuo Yao — 325 — Dibazol ion-selective electrodes

J. M. Fernandez-Romero, M. D. Luque de Castro and M. Valcarcel — 333 — Enzymatic determination of total cholesterol in serum by flow injection analysis

B. Monsarrat, M. Maftouh, G. Meunier, J. Bernadou, J. P. Armand, C. Paoletti and B. Meunier — 341 — Oxidative biotransformation of the antitumour agent elliptinium acetate: structural characterization of its human and rat urinary metabolites

S. J. McIntosh, I. D. Watson, P. W. D. Hughes, A. M. Chambers and M. J. Stewart — 353 — Off-line automated sample preparation—experience in a clinical drug investigation unit

K. Shimada, E. Nagashima, S. Orii and T. Nambara — 361 — New derivatization method using ferrocene reagents for the determination of steroid glucuronides by high-performance liquid chromatography with electrochemical detection

R. J. Yarwood, W. D. Moore and J. H. Collett — 369 — The influence of the ammonium ion on the stability of ethacrynic acid in aqueous solution

H. Ibrahim, T. Boye, M. Wermeille, R. Gurny and P. Buri — 379 — Determination of pilocarpine by HPLC in presence of isopilocarpine and a pH-sensitive polymer in ophthalmic dispersions

S. T. Yang, L. O. Wilken and C. R. Clark — 383 — Liquid chromatographic determination of the pH-dependent degradation of eseroline—hydrolysis product of physostigmine

Short Communications

P. Gonzalez-Hierro, M. M. Velazquez, J. M. Cachaza and L. J. Rodriguez — 395 — Deconvolution of ultraviolet absorption spectra of aqueous Diazepam, Flurazepam and Prazepam in acid and alkaline solutions

M. Takayanagi, S. Goto, T. Fukuda and T. Yashiro — 403 — Determination of human serum ceruloplasmin by measurement of its 3,3′,5,5′-tetramethylbenzidine oxidase activity

M. C. Gutierrez, A. Gomez-Hens and M. Valcarcel — 409 — Kinetic-fluorimetric determination of pilocarpine in ophthalmic solutions

D. G. Muller and V. J. Stella — 415 — An analytical method to determine small quantities of phenytoin in the presence of excess amounts of one of its prodrugs, the monomethane sulphonate salt of 3-(*N*,*N*-dimethylglycyloxymethyl)-5,5-diphenylhydantoin

P. J. Palermo — 423 — High pressure liquid chromatographic determination of dyclonine hydrochloride and its degradation products in cough lozenges and liquids

M. A. Abounassif, E.-R. A. Gad Kariem and H. Y. Aboul-Enein — 431 — High-performance liquid chromatographic determination of clobazam and one of its pharmaceutical formulations

E. J. D. Lee and S. B. Ang — 435 — Measurement of debrisoquine and 4-hydroxydebrisoquine in urine by liquid chromatography

Number 5

Review

D. E. Moore — 441 — Principles and practice of drug photodegradation studies

Analytical Survey

B. V. Fisher and R. Jones — 455 — Chemometrics in pharmaceutical analysis

Full Papers

M. Apfel, L. A. Sternson, A. J. Repta and D. Mills — 469 — Design and synthesis of a theophylline bonded-phase column for HPLC—application to separation of aromatic carboxylic acids

P. Leroy and A. Nicolas — 477 — Determination of atropine in pharmaceutical dosage forms containing vegetal preparations, by high-performance liquid chromatography with U.V. and electrochemical detection

P. van Zoonen, C. Gooijer, N. H. Velthorst, R. W. Frei, J. H. Wolf, J. Gerrits and F. Flentge — 485 — HPLC detection of choline and acetylcholine in serum and urine by an immobilized enzyme reactor followed by chemiluminescence detection

A. Colautti, F. Fontani and V. Maurich — 493 — HPLC determination of papaveraldine and papaverinol in papaverine injection

M. S. Bull and G. R. E. Shume — 501 — Simultaneous determination of fenbendazole, its sulphoxide and sulphone metabolites with the corresponding metabolites of triclabendazole in the plasma of sheep and cattle by high-performance liquid chromatography

Short Communications

S. A. Coran, M. Bambagiotti-Alberti, F. F. Vincieri and G. Moneti — 509 — Rapid monitoring of biologically-active substances in medicinal plants by tandem mass spectrometry—the identification of lignans in *Oenanthe aquatica* Lam

M. Niyaz Khan — 515 — Spectrophotometric determination of alkyl salicylate from the mixture of phenyl and alkyl salicylates by the use of the selective reactivity of phenyl salicylate toward secondary amines

J. C. Berridge and L. A. Broad — 523 — The determination of fluconazole in rodent diet using solid-phase extraction and high-performance liquid chromatography

M. S. Bull and G. R. E. Shume — 527 — A rapid high-performance liquid chromatographic procedure for the determination of triclabendazole and its metabolites in sheep plasma

Number 6

Full Papers

N. Daoud, T. Arvidsson and K.-G. Wahlund — 533 — The determination of lidocaine and its metabolites by a direct injection LC method using large sample volumes of untreated plasma

M. H. Golden and P. H. Zoutendam — 543 — The determination of minoxidil in human serum by high-performance liquid chromatography with amperometric detection

F. G. Lopez, E. L. Mariño and A. Dominguez-Gil — 553 — Determination of bentazepam in plasma by high-performance liquid chromatography

J. A. de Schutter, W. van den Bossche and P. de Moerloose — 559 — Simultaneous effects of amines and alkylsulphonates in reversed-phase ion-pair liquid chromatography—application to the separation of 2-imidazoline drugs

B. Morelli — 577 — Spectrophotometric assay for chloramphenicol and some derivatives in the pure form and in formulations

S. Olajos and D. Sztaniszlav — 585 — Determination of fluperlapine in human plasma by capillary gas chromatography

O. R. Idowu — 591 — Detection and estimation of antimalarial drugs in body fluids—I. Field test for amodiaquine in urine and saliva

Short Communications

R. Karunanithy, S. L. Leung and S. H. Yeo — 597 — Use of high-performance liquid chromatography for quantitative structure–activity relationship studies of a series of potential contraceptive steroidal esters

G. Hanna and C. Lau-Cam — 605 — Determination of ecothiopate iodide by ^{1}H-nuclear magnetic resonance spectroscopy

D. M. Barends, J. C. A. M. Brouwers and A. Hulshoff — 613 — Fast pre-column derivatization of aminoglycosides with 1-fluoro-2,4-dinitrobenzene and its application to pharmaceutical analysis

J. Fidanza and J. J. Aaron — 619 — The analysis of pharmaceutical preparations using solid-surface room temperature photochemical-fluorescence

K. Umprayn, P. E. Kennedy, J. C. Lee, W. Waugh, C. M. Riley and V. J. Stella — 625 — Development of reversed-phase HPLC assays for cyclodisone and clomesone

Number 7

Review

E. G. Salole — 635 — The physicochemical properties of oestradiol

Analytical Survey

M. P. Bailey, B. F. Rocks and C. Riley — 649 — On the use of fluorescent labels in immunoassay

Full Papers

Y. Michotte, M. Moors, D. Deleu, P. Herregodts and G. Ebinger — 659 — Simultaneous determination of levodopa, carbidopa, 3-*O*-methyldopa and dopamine in plasma using high-performance liquid chromatography with electrochemical detection

P. Slégel, G. Vereczkey-Donáth, L. Ladányi and M. Tóth-Lauritz — 665 — Enantiomeric separation of chiral carboxylic acids, as their diastereomeric carboxamides, by thin-layer chromatography

K. Umprayn, V. J. Stella and C. M. Riley — 675 — Stability indicating assay for fetindomide (NSC 373965), a potential prodrug of mitindomide (NSC 284356), employing high-performance liquid chromatography

M. A. Ribick, M. Jemal and A. I. Cohen — 687 — Determination of ethylenediaminetetra-acetic acid in aqueous rinses of detergent-washed rubber stoppers of pharmaceutical vials using solid-phase extraction and capillary gas chromatography

P. Miskolczi, L. Vereczkey and R. Frenkl — 695 — Gas–liquid chromatographic method for the determination of tolperisone in human plasma: pharmacokinetic and comparative bioavailability studies

P. Linares, J. Ruz, M. D. Luque de Castro and M. Valcarcel — 701 — Enzymatic determination of ethanol in saliva by flow injection analysis

M. Cornarakis-Lentzos and P. R. Cowin — 707 — Dilutions of corticosteroid creams and ointments—a stability study

Short Communications

F. Małecki, J. C. Crawhall and R. Staróscik — 717 — Determination of melphalan in the perfusate from patients treated by regional perfusion chemotherapy

R. J. Ruane and I. D. Wilson — 723 — The use of C18 bonded silica in the solid phase extraction of basic drugs—possible role for ionic interactions with residual silanols

H. N. Alkaysi, M. S. Salem and A. A. Badwan — 729 — Reversed-phase high-performance liquid chromatographic analysis of terfenadine

I. W. Wainer, M. C. Alembik and L. J. Fischer — 735 — The determination of (*R*)- and (*S*)-glutethimide and the corresponding 4-hydroxyglutethimide metabolites in human serum and urine using a Pirkle-type HPLC chiral stationary phase

M. P. Bailey, B. F. Rocks and C. Riley — 741 — Theoretical considerations on the fluorescence lifetime of labels for time-resolved fluorescence immunoassay

A. El-Yazigi and A. Al-Humaidan — 747 — Rapid analysis of 5-fluorouracil in plasma or formulations by high-performance liquid chromatography

755 — Book Review

Number 8

Special Issue: Immunoassays

A. F. Fell and J. N. Miller — 757 — Editorial

Full Papers

E. Granström and H. Kindahl — 759 — Quality control of antibodies with special reference to prostaglandins

L. Molin, B. Bergdahl and G. Dahlström — 767 — Problems with the immunoassay of digoxin

C. Hogstrand, P.-E. Olsson and C. Haux — 777 — Partial development of a radioimmunoassay for horse metallothionein

C. A. K. Borrebaeck — 783 — Development of *in vitro* immunization in murine and human hybridoma technology

W. S. Prince, K. J. Harder, S. Saks, B. R. Reed, A. B. Chen and A. J. S. Jones — 793 — ELISA for quantitation of tumor necrosis factor-α in serum

A. M. Johnson — 803 — Nephelometric immunoassay

C. J. Stanley, D. H. Ellis, D. L. Bates and A. Johannsson — 811 — Enzyme-amplified immunoassays

J. L. Boitieux, M. P. Biron and D. Thomas — 821 — Development of a reversible bienzymatic system in immunoenzymatic technique

Short Communication

L. J. Kricka and T. P. Whitehead — 829 — Chemiluminescent and bioluminescent immunoassays